

Xifaxan Xifaxan (Rifaximin) is indicated for the treatment of patients (greater than or equal to 12 years of age) with traveler's diarrhea caused by non-invasive strains of Escherichia coli.
 Do not use rifaximin tablets in patients with diarrhea complicated by fever, blood in the stools or diarrhea due to pathogens other than Escherichia coli.

Member Information

Relationship (Circle)

Name _____

Enrollment / Cardholder ID _____

Group / Plan _____

Birth date _____

Address: _____

City State Zip _____

Subscriber Spouse Dependent

Gender (Circle)

Male Female

AGE: _____

Practitioner Information

Name _____

Agent / Contact Name _____

Specialty _____

Office / Clinic Name _____

Address _____

City State Zip _____

Phone () - _____

Fax () - _____

Email _____ @ _____

Is the patient greater than or equal to 12 years of age? Y N

Does the patient have fever? Y N

Does the patient have blood in the stools? Y N

Does the patient have traveler's diarrhea caused by Escherichia coli? Y N

Requested Drug / Pharmaceutical

Xifaxan – Rifaximin

Strength / Dose

Duration of Therapy (Expected)

Condition / Diagnosis Related - Reason for Request

Type of Request (Circle)

One-Time On-Going

Clinical Drug / Lab History Pertinent to Request

Contact Pharmacy? Y N

Pharmacy Phone Number

Formulary Alternative(s) Attempted YES NO (Circle)

Practitioner Signature